

LETTERS TO THE EDITOR

doi:10.1093/occmed/kqv184

Burnout in firefighters: a word on methodology

Dear Sir,

Katsavouni *et al.* [1] examined burnout, post-traumatic stress disorder and work-related injuries in Greek firefighters. The authors found that 20% of the firefighters in their sample met criteria for burnout as treated as a nosological entity. To receive a diagnosis of burnout, scores on the emotional exhaustion, depersonalization and personal accomplishment subscales of the Maslach Burnout Inventory had to be above or below certain thresholds. At least three methodological problems affect the study.

First, there are currently no diagnostic criteria for burnout [2], neither in the DSM-5, nor in the ICD-10. Second, although for any study, a critic could assert that a key variable was omitted, we believe that one extremely important variable was indeed omitted from this study, namely, depression. While Katsavouni *et al.* [1] defined burnout as ‘a long-term reaction to occupational stress’, there is a great deal of evidence that depressive symptoms and disorders develop in response to exposure to occupational stressors [3–8]. Moreover, evidence that depression subsumes burnout has been mounting [9–12]. Ahola *et al.* [9] and Bianchi *et al.* [12] found that burnout and depressive symptoms increase or decline commensurately over time. Ahola *et al.* [9] concluded that ‘burnout could be used as an equivalent to depressive symptoms in work life’ (p. 35). Bianchi *et al.* [12] suggested that the burnout-depression distinction is ‘symptomatic of different scientific traditions, rather than of different pathological phenomena’ (p. 1010). Thus, there is no need to multiply constructs when the long-studied construct depression can serve researchers well. It is noteworthy that several firefighters who experienced traumatic events (e.g. death of a child) mentioned experiencing *depression* regarding their feelings of responsibility. Judging from Table 2, none mentioned burnout.

Third, the authors did not control for relevant non-occupational factors such as stressors occurring outside of work. To be able to conclude that the symptoms under examination were specifically related to *work*, it is necessary to take into account the stressors experienced in other life domains. For instance, burnout is associated with personal/stressful life events (e.g. a loved one suffering a major illness) [11,13]. In our view, research assessing dispositional (e.g. personality variables) and situational (both occupational and non-occupational) factors is needed in occupational medicine so that clear conclusions can be

drawn regarding the influence of work-related exposures in the emergence of burnout/depression.

Irvin Sam Schonfeld

Department of Psychology,
The City College and the Graduate Center of the City,
University of New York,
New York, NY, USA
e-mail: ischonfeld@ccny.cuny.edu

Renzo Bianchi

Institute of Work and Organizational Psychology,
University of Neuchâtel,
Neuchâtel, Switzerland

References

1. Katsavouni F, Bebetos E, Malliou P, Beneka, A. The relationship between burnout, PTSD symptoms and injuries in firefighters. *Occup Med (Lond)* 2016;**66**:32–37.
2. Bianchi R, Schonfeld IS, Laurent E. Burnout-depression overlap: a review. *Clin Psychol Rev* 2015;**36**:28–41.
3. Bonde JP. Psychosocial factors at work and risk of depression: a systematic review of the epidemiological evidence. *Occup Environ Med* 2008;**65**:438–445.
4. Netterstrom B, Conrad N, Bech P *et al.* The relation between work-related psychosocial factors and the development of depression. *Epidemiol Rev* 2008;**30**:118–132.
5. Rydmark I, Wahlberg K, Ghatan PH *et al.* Neuroendocrine, cognitive and structural imaging characteristics of women on longterm sickleave with job stress-induced depression. *Biol Psychiatry* 2006;**60**:867–873.
6. Tennant C. Work-related stress and depressive disorders. *J Psychosom Res* 2001;**51**:697–704.
7. Theorell T, Hammarström A, Aronsson G *et al.* A systematic review including meta-analysis of work environment and depressive symptoms. *BMC Public Health* 2015;**15**, 738.
8. Wang J. Work stress as a risk factor for major depressive episode(s). *Psychol Med* 2005;**35**:865–871.
9. Ahola K, Honkonen T, Isometsä E *et al.* The relationship between job-related burnout and depressive disorders—results from the Finnish Health 2000 Study. *J Affect Disord* 2005;**88**:55–62.
10. Bianchi R, Schonfeld IS, Laurent E. Is burnout a depressive disorder? A re-examination with special focus on atypical depression. *Int J Stress Manag* 2014;**21**:307–324.
11. Schonfeld, IS, Bianchi R. Burnout and depression: two entities or one. *J Clin Psychol* 2015. doi:10.1002/jclp.22229.
12. Bianchi R, Schonfeld IS, Laurent E. Is burnout separable from depression in cluster analysis? A longitudinal study. *Soc Psychiatry Psychiatr Epidemiol* 2015;**50**:1005–1011.
13. Dyrbye LN, Thomas MR, Huntington JL *et al.* Personal life events and medical student burnout: a multicenter study. *Acad Med* 2006;**81**:374–384.